

(19)



Europäisches Patentamt  
European Patent Office  
Office européen des brevets



(11)

**EP 1 382 337 A1**

(12)

**EUROPEAN PATENT APPLICATION**

(43) Date of publication:

**21.01.2004 Bulletin 2004/04**

(51) Int Cl.7: **A61K 31/445**, A61K 9/20,  
A61K 31/4525

(21) Application number: **03078259.3**

(22) Date of filing: **19.07.1996**

(84) Designated Contracting States:

**AT BE CH DE DK ES FI FR GB GR IE IT LI LU MC  
NL PT SE**

Designated Extension States:  
**SI**

(30) Priority: **20.07.1995 GB 9514842**

(62) Document number(s) of the earlier application(s) in  
accordance with Art. 76 EPC:  
**96926388.8 / 0 839 039**

(71) Applicant: **SMITHKLINE BEECHAM PLC**  
**Brentford, Middlesex TW8 9GS (GB)**

(72) Inventors:

- **Leonard, Graham S. c/o GlaxoSmithKline  
Harlow Essex CM19 5AW (GB)**
- **Elder, David P. c/o GlaxoSmithKline  
Ware Hertfordshire SG12 0DP (GB)**

(74) Representative: **Knight, Lucie Viktoria**  
**GlaxoSmithKline,**  
**C.I.P., CN925.1.980, Great West Road**  
**Brentford, Middlesex TW8 9GS (GB)**

Remarks:

This application was filed on 15/10/2003 as a  
divisional application to the application mentioned  
under INID code 62.

(54) **Novel formulation containing paroxetine**

(57) Controlled release and delayed release formulations which are adapted or intended for oral administration and which contain paroxetine hydrochloride, such as a formulation which is a system for the controlled and delayed release of paroxetine hydrochloride, having (a) a deposit-core comprising 28.61 mg paroxetine hydrochloride, 18.7 mg methocel K4M, 79.14 mg lactose monohydrate, 2.50 mg polyvinylpyrrolidone, 1.25 mg magnesium stearate, and 0.50 mg Syloid 244,

and (b) a support-platform applied to said deposit-core comprising 15.04 mg Compritol 888, 30.50 mg lactose monohydrate, 4.00 mg polyvinylpyrrolidone, 0.80 mg magnesium stearate, 29.32 mg methocel E5, 0.32 mg Syloid 244, and 0.02 mg iron oxide; and (c) an enteric coating comprising 13.27 mg Eudragit, 3.31 mg talc, and 1.33 mg triethyl citrate, may be used to treat a variety of disorders including depression.

**EP 1 382 337 A1**

## Description

[0001] The present invention relates to a novel formulation containing paroxetine or a pharmaceutically acceptable salt thereof, and to its use in the treatment and/or prophylaxis of certain disorders.

[0002] US Patent No 4,007,196 describes *inter alia* a compound which is commonly known as paroxetine. This compound is a Selective Serotonin Reuptake Inhibitor (SSRI) and is currently marketed world-wide for the treatment and/or prophylaxis of depression.

[0003] The current formulation which is the only marketed formulation of paroxetine hydrochloride is a swallow tablet.

[0004] It has now been surprisingly found that controlled release and delayed release formulations containing paroxetine give rise to an unexpected reduction in the side effects associated with swallow tablets.

[0005] Accordingly, the present invention provides a controlled release or delayed release formulation containing paroxetine or a pharmaceutically acceptable salt thereof.

[0006] A further aspect of the invention provides a controlled release or delayed release formulation containing an SSRI. Examples of SSRIs other than paroxetine include fluoxetine (US Patent No. 4,314,081), fluvoxamine (US Patent No. 4,085,225), and sertraline (US Patent No. 4,536,518).

[0007] By controlled release is meant any formulation technique wherein release of the active substance from the dosage form is modified to occur at a slower rate than that from an immediate release product, such as a conventional swallow tablet or capsule.

[0008] By delayed release is meant any formulation technique wherein release of the active substance from the dosage form is modified to occur at a later time than that from a conventional immediate release product. The subsequent release of active substance from a delayed release formulation may also be controlled as defined above.

[0009] Examples of controlled release formulations which are suitable for incorporating paroxetine and other SSRIs are described in:

Sustained Release Medications, Chemical Technology Review No. 177. Ed. J.C. Johnson. Noyes Data Corporation 1980.

Controlled Drug Delivery, Fundamentals and Applications, 2nd Edition. Eds. J.R. Robinson, V.H.L. Lee. Marcel Dekkes Inc. New York 1987.

[0010] Examples of delayed release formulations which are suitable for incorporating paroxetine and other SSRIs are described in:

[0011] Remington's Pharmaceutical Sciences 16th Edition, Mack Publishing Company 1980, Ed. A. Osol.

[0012] Such controlled release formulations are preferably formulated in a manner such that release of active substance such as paroxetine is effected predominantly during the passage through the stomach and the small intestine, and delayed release formulations are preferably formulated such that release of active substance such as paroxetine is avoided in the stomach and is effected predominantly during passage through the small intestine

[0013] Said formulations are preferably formulated such that the release of the active substance is predominantly 1½ to 3 hours post ingestion.

[0014] The small intestine is suitably the duodenum, the ileum or the jejunum.

[0015] Patients who benefit most from the formulations of the present invention are those who are known to suffer from nausea upon oral administration using swallow tablets.

[0016] Preferred formulations are ultimately enteric coated tablets or caplets, wax or polymer coated tablets or caplets or time-release matrices, or combinations thereof.

[0017] Particularly preferred formulations are described in US Patent No. 5,102,666.

[0018] Thus, a particular aspect of the invention provides a polymeric controlled release composition comprising a reaction complex formed by the interaction of (1) a calcium polycarbophil component which is a water-swellaable, but water insoluble, fibrous cross-linked carboxy-functional polymer, said polymer containing (a) a plurality of repeating units of which at least about 80% contain at least one carboxyl functionality, and (b) about 0.05 to about 1.5% cross-linking agent substantially free from polyalkenyl polyether, said percentages being based upon the weights of unpolymerised repeating unit and cross-linking agent, respectively, with (2) water, in the presence of an active agent selected from the group consisting of SSRIs such as paroxetine. The amount of calcium polycarbophil present is from about 0.1 to about 99% by weight, for example about 10%. The amount of active agent present is from about 0.0001 to about 65% by weight, for example between about 5 and 20%. The amount of water present is from about 5 to about 200% by weight, for example between about 5 and 10%. The interaction is carried out at a pH of between about 3 and about 10, for example about 6 to 7. The calcium polycarbophil is originally present in the form of a calcium salt containing from about 5 to about 25% calcium.

[0019] Further particularly preferred formulations are described in US Patent No. 5,422,123.

[0020] Thus, a further particular aspect of the invention provides a system for the controlled release of an active

substance which is an SSRI such as paroxetine, comprising (a) a deposit-core comprising an effective amount of the active substance and having defined geometric form, and (b) a support-platform applied to said deposit-core, wherein said deposit-core contains at least the active substance, and at least one member selected from the group consisting of (1) a polymeric material which swells on contact with water or aqueous liquids and a gellable polymeric material wherein the ratio of the said swellable polymeric material to said gellable polymeric material is in the range 1:9 to 9:1, and (2) a single polymeric material having both swelling and gelling properties, and wherein the support-platform is an elastic support, applied to said deposit-core so that it partially covers the surface of the deposit-core and follows changes due to hydration of the deposit-core and is slowly soluble and/or slowly gellable in aqueous fluids. The support-platform may comprise polymers such as hydroxypropylmethylcellulose, plasticizers such as a glyceride, binders such as polyvinylpyrrolidone, hydrophilic agents such as lactose and silica, and/or hydrophobic agents such as magnesium stearate and glycerides. The polymer(s) typically make up 30 to 90% by weight of the support-platform, for example about 35 to 40%. Plasticizer may make up at least 2% by weight of the support-platform, for example about 15 to 20%. Binder(s), hydrophilic agent(s) and hydrophobic agent(s) typically total up to about 50% by weight of the support-platform, for example about 40 to 50%.

[0021] Paroxetine used in the present invention is suitably in the form of the free base or a pharmaceutically acceptable salt thereof. Preferably, paroxetine is suitably in the form of the hydrochloride hemihydrate.

[0022] Paroxetine hydrochloride hemihydrate may be prepared according to the procedures generally outlined in US Patent 4,721,723..

[0023] Paroxetine in the form of a controlled release or delayed release formulation can be used to treat and prevent the following disorders:

- Alcoholism
- Anxiety
- Depression
- Obsessive Compulsive Disorder
- Panic Disorder
- Chronic Pain
- Obesity
- Senile Dementia
- Migraine
- Bulimia
- Anorexia
- Social Phobia
- Pre-Menstrual Syndrome (PMS)
- Adolescent Depression
- Trichotillomania
- Dysthymia
- Substance Abuse

[0024] These disorders are herein after referred to as "the disorders".

[0025] The present invention provides a method of treating and/or preventing the disorders by administering an effective and/or a prophylactic amount of a controlled release or delayed release formulation containing paroxetine or a pharmaceutically acceptable salt thereof, to a sufferer in need thereof.

[0026] The present invention further provides the use of a controlled release or delayed release formulation containing paroxetine or a pharmaceutically acceptable salt thereof in the manufacture of a medicament, for treating and/or preventing the disorders.

[0027] The present invention also provides a pharmaceutical composition for use in the treatment and/or prevention of the disorders which comprises a controlled release or delayed release formulation containing paroxetine or a pharmaceutically acceptable salt thereof.

[0028] The following examples illustrate the present invention.

#### Example 1 (Hydrophilic Matrix)

[0029]

Intragranular	% w/w
Paroxetine Hydrochloride	11.45

# EP 1 382 337 A1

(continued)

Intragranular	% w/w
Methocel E5	1.25
Lactose	12.3
Extragranular	
Methocel K100LV	30.0
Lactose	44.0
Magnesium Stearate	1.0
TOTAL	100.0

## Example 2 (Hydrophilic Matrix)

[0030]

Intragranular	% w/w
Paroxetine Hydrochloride	11.45
Methocel E5	1.25
Lactose	12.3
Extragranular	
Methocel K100LV	27.5
Methocel K4M	7.5
Lactose	39.0
Magnesium Stearate	1.0
TOTAL	100.0

## Example 3 (pH Sensitive Coat on Immediate Release Core)

[0031]

Tablet Core	%w/w
Paroxetine Hydrochloride	11.45
Lactose	64.05
Microcrystalline Cellulose	20.0
Sodium Starch Glycollate	4.0
Magnesium Stearate	0.5
TOTAL	100.0

Tablet Coating (apply approximately 6-10% of tablet core weight)	%w/w
Hydroxypropylmethylcellulose Phthalate	90.0
Triacetin	10.0

## Example 4 (pH Sensitive Coat on Immediate Release Core)

[0032] Tablet Core as in Example 3

Tablet Coating (apply approximately 6-10% of tablet core weight)	%w/w
Cellulose Acetate Phthalate	90.0
Diethyl Phthalate	10.0

## EP 1 382 337 A1

### Example 5 (Controlled Release Coating on Immediate Release Core)

[0033] Tablet Core as in Example 3

Tablet Coating (apply approximately 5-12% of tablet core weight)	%w/w
Eudragit RS 100	86.0
Dibutyl Phthalate	10.0
Talc	4.0
FD&C Yellow No. 6	0.01

### Example 6 (pH Sensitive Coat on Controlled Release Core.)

[0034] Tablet Core as in Example 3

[0035] Tablet Coating as in Example 3

### Example 7 (Encapsulated Controlled Release Coated Beads)

[0036]

Pellet	%w/w (approx)
Non Pareil Seed	30
Paroxetine Hydrochloride	40
Gelatin	8
Lactose	20
Talc	2

Coating	%w/w
Glycerylmonostearate	36.6
Glyceroldistearate	53.4
White Wax	10.0

### Example 8 (Controlled release bilayer tablet)

[0037]

Active Layer		
Component	mg/tablet	Function
Paroxetine Hydrochloride	22.89*	Active
Methocel K4M	15.00	Hydrogel polymer
Lactose monohydrate	62.0	Hydrophilic agent
Polyvinylpyrrolidone	3.0	Binder
Magnesium stearate	1.0	Hydrophobic agent
Syloid 244	1.0	Hydrophilic agent
Support platform		
Component	mg/tablet	Function
Compritol 888	15.04	Plasticizer
Lactose monohydrate	29.32	Hydrophilic agent
Polyvinylpyrrolidone	4.0	Binder

\*Equivalent to 20mg paroxetine as free base.

# EP 1 382 337 A1

(continued)

Support platform		
Component	mg/tablet	Function
Magnesium stearate	1.52	Hydrophobic agent
Methocel E5	29.32	Hydrogel polymer
Iron oxide	0.08	Colourant
Total tablet weight	184.89mg	

[0038] The powder blend for each layer was wet granulated in a high shear mixer/granulator and dried in a fluid bed drier. The bilayer tablets were compressed on a Manesty triple layer press.

Example 9 (Enteric coated calcium polycarbophil formulation)

[0039]

Core		
Component	mg/ tablet	Function
Paroxetine Hydrochloride	22.89*	Active
Calcium polycarbophil	20.00	Matrix
Lactose anhydrous	146.11	Hydrophilic agent/diluent
Polyvinylpyrrolidone	10.0	Binder
Magnesium stearate	1.0	Hydrophobic agent/lubricant
Water**	0.024	Granulating liquid
Enteric coat		
Component	mg/ tablet	Function
Eudragit	22.19	Polymer
Talc	1.53	Lubricant
Triethyl citrate	1.00	Plasticizer
Water**	24.6	Diluent
Film coat		
Opadry pink	10.5	Film coat
Water**	94.5	Diluent
Polish coat		
Opadry clear	0.750	
Water**	29.3	Diluent

\*Equivalent to 20mg paroxetine as free base.

\*\*Removed during processing.

[0040] The core constituents were wet granulated in a high shear mixer/granulator, and dried in a fluid bed drier. The magnesium stearate was then added and the mixture processed in a low shear mixer. The mix was then compressed on a B type rotary tablet press. Coating was carried out using an Accela cota.

# EP 1 382 337 A1

Example 10 (Controlled release bilayer tablet)

[0041]

Active Layer		
Component	mg/tablet	Function
Paroxetine Hydrochloride	22.89*	Active
Methocel K4M	20.00	Hydrogel polymer
Lactose monohydrate	60.0	Hydrophilic agent
Polyvinylpyrrolidone	5.0	Binder
Magnesium stearate	1.0	Hydrophobic agent
Syloid 244	1.0	Hydrophilic agent
Support platform		
Component	mg/tablet	Function
Compritol 888	14.72	Plasticizer
Lactose monohydrate	30.60	Hydrophilic agent
Polyvinylpyrrolidone	2.80	Binder
Magnesium stearate	0.80	Hydrophobic agent
Methocel E5	30.60	Hydrogel polymer
Syloid 244	0.40	Hydrophilic agent
Iron oxide	0.08	Colourant
Total tablet weight	189.89mg	

\*Equivalent to 20mg paroxetine as free base.

[0042] The process was as described in Example 8.

Example 11 (Controlled release bilayer tablet)

[0043]

Active Layer		
Component	mg/tablet	Function
Paroxetine Hydrochloride	22.89*	Active
Methocel K4M	15.00	Hydrogel polymer
Lactose monohydrate	63.31	Hydrophilic agent
Polyvinylpyrrolidone	2.0	Binder
Magnesium stearate	1.0	Hydrophobic agent
Syloid 244	0.40	Hydrophilic agent
Support platform - as in Example 10.		
Total tablet weight	184.60mg	

\*Equivalent to 20mg paroxetine as free base.

[0044] The process was as described in Example 8.

## EP 1 382 337 A1

Example 12 (Enteric coated controlled release bilayer tablet)

[0045]

Active Layer		
Component	mg/tablet	Function
Paroxetine Hydrochloride	28.61*	Active
Methocel K4M	18.75	Hydrogel polymer
Lactose monohydrate	79.14	Hydrophilic agent
Polyvinylpyrrolidone	2.50	Binder
Magnesium stearate	1.25	Hydrophobic agent
Syloid 244	0.50	Hydrophilic agent
Support platform		
Component	mg/tablet	Function
Compritol 888	15.04	Plasticizer
Lactose monohydrate	30.50	Hydrophilic agent
Polyvinylpyrrolidone	4.00	Binder
Magnesium stearate	0.80	Hydrophobic agent
Methocel E5	29.32	Hydrogel polymer
Syloid 244	0.32	Hydrophilic agent
Iron oxide	0.02	Colourant
Enteric coating		
Component	mg/tablet	Function
Eudragit	13.27	Polymer
Talc	3.31	Lubricant
Triethyl citrate	1.33	Plasticizer
Water**	36.25	Diluent
Total tablet weight	228.66mg	

\*Equivalent to 25mg paroxetine as free base.

\*\*Removed during processing.

[0046] The process was as described in Example 9.

Example 13

### GI tolerance study

[0047] The design of the study is outlined below

Subjects: Normal healthy volunteers  
 Design: Parallel group, placebo controlled, double blind  
 Treatment: (a) Placebo, (b) Immediate release paroxetine, (c) Example 8 formulation, (d) Example 8 formulation with enteric coating.  
 Dosage: 30 mg once daily for 3 days  
 Number of subjects: 452 evaluable (488 randomised, 485 evaluable)

[0048] The study was conducted to compare the incidence, severity and duration of nausea and vomiting, and diarrhoea (theoretically if the controlled release formulations slow down absorption of paroxetine then, as paroxetine is known to be prokinetic to the GI tract there may be an increased incidence).

[0049] Adverse experiences (AE) information was assessed each morning at the time of dosing and again 24 hours following the last dose. Investigators and subjects were given diary cards detailing how to classify severity of AEs in order to standardise as much as possible across all 6 centres.



## EP 1 382 337 A1

[0050] Of the 485 evaluable subjects, 18 (3.7%) withdrew, 17 because of adverse events. Subjects with nausea/vomiting on the day of withdrawal were more common on (b) than either of (c) and (d).

[0051] The incidence of nausea/vomiting and diarrhoea is shown in the table below:

	(b)	(c)	(d)	Placebo
Incidence of nausea	59%	49%	39%	13%
Incidence of diarrhoea	15%	21%	20%	7%

[0052] The incidence of nausea was increased for both (b) and placebo compared to the expected rates of approximately 25% and 5% respectively for volunteers at these dosages for 3 days duration. The overall incidence of nausea was less on (c) and (d) than on (b). The severity of nausea was also decreased as shown in the next table.

Nausea severity	(b)	(c)	(d)	Placebo
None	50 (41%)	63 (52%)	74 (61 %)	104 (87%)
Mild	45 (37%)	40 (33%)	30 (25%)	16 (13%)
Moderate	21 (17%)	17 (14%)	15 (12%)	0 (0%)
Severe	6 (5%)	1 (1%)	3 (2%)	0 (0%)

[0053] Severity of diarrhoea is reported in the table below:

Severity of diarrhoea	(b)	(c)	(d)	Placebo
None	104 (85%)	95 (79%)	97 (80%)	112 (93%)
Mild	16 (13%)	16 (13%)	16 (13%)	8 (7%)
Moderate	1 (1%)	8 (7%)	9 (7%)	0 (0%)
Severe	1 (1%)	2 (2%)	0 (0%)	0 (0%)

[0054] In conclusion, there appears to be a trend for (c) to reduce the incidence of nausea and the dropout rate due to adverse events in comparison to (b), but analysis of the results was complicated by a statistically significant treatment-by-centre difference. (d) shows a halving in the dropout rate and a fall in incidence of nausea of 20% (a proportional fall of 33%). In addition there is a reduction in severity of nausea of those individuals who report nausea on (c) and (d). There is an increase in incidence of diarrhoea on both of (c) and (d) in relation to (b), but this is confined to an increase in the number of individuals reporting moderate diarrhoea and there is no increase in those with severe diarrhoea.

### Claims

1. A controlled release and delayed release formulation which is adapted or intended for oral administration and which contains paroxetine hydrochloride.
2. A controlled release and delayed release formulation according to claim 1 wherein paroxetine hydrochloride is in the form of paroxetine hydrochloride hemihydrate.
3. A tablet formulation which is a system for the controlled and delayed release of paroxetine hydrochloride, having (a) a deposit-core comprising 28.61 mg paroxetine hydrochloride, 18.75 mg methocel K4M, 79.14 mg lactose monohydrate, 2.50 mg polyvinylpyrrolidone, 1.25 mg magnesium stearate, and 0.50 mg Syloid 244; and (b) a support-platform applied to said deposit-core comprising 15.04 mg Compritol 888, 30.50 mg lactose monohydrate, 4.00 mg polyvinylpyrrolidone, 0.80 mg magnesium stearate, 29.32 mg methocel E5, 0.32 mg Syloid 244, and 0.02 mg iron oxide; and (c) an enteric coating comprising 13.27 mg Eudragit, 3.31 mg talc, and 1.33 mg triethyl citrate.



European Patent  
Office

## EUROPEAN SEARCH REPORT

Application Number  
EP 03 07 8259

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int.Cl.7)
A	WO 95 15155 A (SMITHKLINE BEECHAM) 8 June 1995 (1995-06-08) * the whole document *	1-3	A61K31/445 A61K9/20 A61K31/4525
A,D	EP 0 432 607 A (JAGOTEC AG) 19 June 1991 (1991-06-19) * the whole document *	1-3	
A,D	WO 92 03124 A (ORAMED) 5 March 1992 (1992-03-05) * the whole document *	1-3	
A,P	CA 2 143 070 A (P.MODI) 23 August 1995 (1995-08-23) * claims *	1-3	
			TECHNICAL FIELDS SEARCHED (Int.Cl.7)
			A61K
The present search report has been drawn up for all claims			
Place of search THE HAGUE		Date of completion of the search 25 November 2003	Examiner Scarponi, U
<p>CATEGORY OF CITED DOCUMENTS</p> <p>X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document</p> <p>T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons &amp; : member of the same patent family, corresponding document</p>			

EPO FORM 1503 03.82 (P04C01)

**ANNEX TO THE EUROPEAN SEARCH REPORT  
ON EUROPEAN PATENT APPLICATION NO.**

EP 03 07 8259

This annex lists the patent family members relating to the patent documents cited in the above-mentioned European search report.  
The members are as contained in the European Patent Office EDP file on  
The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

25-11-2003

Patent document cited in search report		Publication date		Patent family member(s)	Publication date
WO 9515155	A	08-06-1995	IT	MI932540 A1	05-06-1995
			AU	693144 B2	25-06-1998
			AU	1219895 A	19-06-1995
			CA	2177721 A1	08-06-1995
			CN	1145586 A	19-03-1997
			WO	9515155 A1	08-06-1995
			EP	0804168 A1	05-11-1997
			JP	9505818 T	10-06-1997
			NZ	277238 A	27-04-1998
			ZA	9409567 A	10-10-1995
EP 432607	A	19-06-1991	IT	1237904 B	18-06-1993
			AT	135906 T	15-04-1996
			CA	2031393 A1	15-06-1991
			DE	69026215 D1	02-05-1996
			DE	69026215 T2	22-08-1996
			DK	432607 T3	29-04-1996
			EP	0432607 A1	19-06-1991
			ES	2085316 T3	01-06-1996
			GR	3020404 T3	30-09-1996
			JP	2907557 B2	21-06-1999
			JP	6172162 A	21-06-1994
			US	5422123 A	06-06-1995
WO 9203124	A	05-03-1992	US	5110605 A	05-05-1992
			US	5102666 A	07-04-1992
			US	5686094 A	11-11-1997
			AT	137404 T	15-05-1996
			AU	8444291 A	17-03-1992
			DE	69119217 D1	05-06-1996
			DE	69119217 T2	31-10-1996
			DK	497956 T3	12-08-1996
			EP	0497956 A1	12-08-1992
			ES	2088500 T3	16-08-1996
			GR	3020502 T3	31-10-1996
			JP	5502894 T	20-05-1993
			JP	3064417 B2	12-07-2000
			WO	9203124 A1	05-03-1992
CA 2143070	A	23-08-1995	CA	2143070 A1	23-08-1995

EPO FORM P055

For more details about this annex : see Official Journal of the European Patent Office, No. 12/82